

## Original Paper

# The Protective Role of Vitamins (C&E), Selenium, Silymarin and Rehydran-N against Lead Toxicity under Heat Stress Conditions

Shalan MG\*, Abd Ali WDh\*\*, El-Batanony MH\*\*\*

### ABSTRACT

This study was done to investigate the use of vitamins C & E, selenium, silymarin and rehydration solution to ameliorate lead toxicity under heat stress conditions. Male albino rats were subdivided into four groups: the first was a control group, the second was exposed to heat stress ( $40 \pm 2$  °C) in a closed and controlled chamber, the third received 25 mg/100 g body weight lead acetate day by day and exposed to the same heat stress conditions, the fourth was exposed to the same lead and hyperthermia conditions and supplemented three times per week with 1 mg/100 g body weight of each of vitamins C & E, silymarin, and 0.01mg/100g B.W. of selenium, and a daily drink of rehydran-n solution. Blood samples were collected after 25 days of treatment. Lead was found to induce significant elevations in blood glucose, total protein, cholesterol, lead, ALT, AST, GGT, ALP and LDH levels under heat stress conditions. Hyperthermia induced apoptotic DNA fragmentation, which was aggravated under lead intoxication. A reduction in body weight was observed in heat stressed groups. Hepatomegaly was observed in heat stressed animals, which was aggravated with lead intake. Under heat stress condition, randomly scattered hepatocytes showed acidophilic and apoptotic changes. Under heat and lead exposure, these changes were enhanced and showed midzonal distribution, in addition to marked periportal microvesicular steatosis. Treatment of rats with vitamins C & E, silymarin, selenium, and rehydran-n resulted in marked improvement in the biochemical, molecular, physiological, and histopathological parameters.

**Key Words:** Rats, Lead, Hyperthermia, Pathology, Apoptosis, Steatosis

### Introduction

Usage of lead and its compounds spread in modern industry.<sup>1</sup> Lead exerts its toxic effects by enhancing peroxidative damage to membranes.<sup>2</sup> Investigations suggest that effects of lead may be due to its interference with calcium in the activation of protein kinase C (PKC) and or through production of reactive oxygen species (ROS).<sup>3</sup> Lead induces DNA strand breaks<sup>4,5</sup> that induce apoptotic DNA fragments reversibly with time.<sup>6</sup> Honchel et al showed that lead enhances lipopolysaccharide and tumor necrosis factor-induced liver injury.<sup>7</sup> Kupffer cells promote lead induced hepatocyte apoptosis via oxidative stress.<sup>8</sup>

Heat stress was reported to induce alterations in hematological,<sup>9</sup> physiological,<sup>10,11</sup> and biochemical<sup>12</sup> parameters. Heat stress exaggerates the toxic effect of lead on different body systems.<sup>13</sup>

Vitamin C has been reported as a chelating agent in treatment of lead toxicity.<sup>14</sup> It reduces the possibility of lead interacting with critical biomolecules and preventing its toxicity.<sup>15</sup> Vitamin E is an antioxidant that helps in capturing reactive oxygen and free radicals produced by toxins.<sup>16</sup> Silymarin is a mixture of antioxidants excreted from *Silybum marianum*. It serves as a free radical scavenger

\*(Author for correspondence): Al Arish Faculty of Education, Suez Canal University, Biological and Geological Department, North Sinai, Egypt. e-mail: mohamedshalan@yahoo.com \*\*High Health Institute, Egdabia, Libya \*\*\*National Liver Institute, Menoufiya University, Histopathology Department, Egypt

and a membrane stabilizer that prevents lipoperoxidation and the associated cell damage.<sup>17</sup> It has a protective effect against experimental hepatotoxicity by regulating liver cell physiology and improving the performance of hepatic enzymes and bile production.<sup>18,19</sup> Vitamin C and silymarin supplements have been used effectively in amelioration of lead toxicity.

The present study aimed to evaluate the effect of using vitamins (C & E), silymarin, selenium and rehydration solution supplements as a protective measures against lead toxicity under heat stress conditions.

## Materials and Methods

### Animals:

Twenty (20) male albino rats (*Rattus norvegicus*), purchased from animal house of Garuonis University, Libya, 10 weeks old, weighing about  $120 \pm 10$  g were used as experimental animals. Animals were housed in groups in plastic cages and were maintained under standard controlled conditions. Laboratory balanced diet and water were initially provided.

### Experimental design:

Animals were segregated into four groups:

- 1 Normal controls.
- 2 Heat stressed group: they were exposed to  $40 \pm 2^\circ\text{C}$  in a closed and controlled chamber and water ad-libitum.
- 3 Lead exposed group (25 mg/100g body weight day by day) under the same heat stress conditions ( $40 \pm 2^\circ\text{C}$ ) and water ad-libitum.
- 4 Lead exposed group (25 mg/100g body weight day by day) under the same heat stress conditions ( $40 \pm 2^\circ\text{C}$ ) and supplemented with vitamin C (1mg/100g body weight), vitamin E (1mg/100g body weight), selenium (0.01 mg/100g body weight) and silymarin (1mg/100 g body weight) by gastric tube 3 times/week and daily free drink rehydran-n solution (4g glucose anhydrous, 0.51g tri-sodium citrate anhydrous, 0.70g sodium chloride and 0.30g potassium chloride in 200 ml distilled water).

### Duration of administration:

Animals of different groups were anaesthetized and rapidly dissected after 25 days of treatment.

### Collection of serum samples:

Blood samples were collected from the abdominal vein

in glass centrifuge tubes, then centrifuged for 15 min. at 1000 x g. Sera were separated and stored at  $-30^\circ\text{C}$  in deep freezer till further biochemical measurements.

### Preparation of tissues for microscopical and gel examinations:

After animal dissection, liver was removed, blotted on filter paper, and weighed. Representative specimens were labeled by code numbers, and immediately placed in 10% formalin. Portions of 10 mg were taken immediately for gel examinations and the remaining portions were stored at  $-30^\circ\text{C}$ .

### Histopathological methods:

The specimens were processed to paraffin as per standard procedure. Five-micron thick histological sections were prepared, and stained with hematoxyline and eosin. The stained sections were examined blindly (without knowing the code number key) under light microscopy. Microscopic examinations of liver sections were done systematically with comments on architecture; portal, periportal, or lobular inflammatory cell infiltrate and its degree; liver cell necrotic or degenerative changes; bile ducts and vascular changes.

### Gel preparation and electrophoresis of lysate tissue:

Gels were prepared with 1.8 % electrophoretic grade agarose (BRL). The agarose was boiled in tris-borate EDTA buffer (1 x TBE buffer; 89 mM tris, 89 mM boric acid, 2 mM EDTA, pH 8.3). 0.5  $\mu\text{g}/\text{ml}$  ethidium bromide was added to gel at  $40^\circ\text{C}$ . Gels were poured and allowed to solidify at room temperature for 1h before samples were loaded. 10 mg hepatic tissue was squeezed and lysed in 200  $\mu\text{l}$  lysing buffer (50 mM NaCl, 1mM Na<sub>2</sub>EDTA, 0.5% SDS, pH 8.3) for at least 30 min.

For electrophoretic pattern of nucleic acids of tissue lysate, 20 $\mu\text{l}$  of lysate hepatic cells was loaded in well, 5  $\mu\text{l}$  6 x loading buffer was on the lysing tissue. Electrophoresis was performed for 2 hours at 50 V in gel buffer (1x TBE buffer). Gel was photographed using a Polaroid camera while the DNA and RNA was visualized using a 312 nm UV transilluminator.

### Nucleic acids extraction and molecular assessment for apoptosis:

Nucleic acids extraction was based on salting out extraction method,<sup>20</sup> whereas protein was precipitated by saturated solution of NaCl (5M). 10mg hepatic tissue

was squeezed in eppendorf tube and was lysed by 600 µl lysing buffer (50 mM NaCl, 1mM Na<sub>2</sub> EDTA, 0.5 % SDS, pH 8.3), and was shaken gently. The mixture was kept overnight at 37°C. For protein precipitation, an amount of 200 µl of saturated NaCl was added to the samples and then shaken gently, and centrifuged at 12,000 rpm for 10 min. The supernatant was transferred to new eppendorf tube and the DNA was precipitated by 700-µl cold iso-propanol. The mixture was inverted several times till fine fibres of nucleic acids appeared, and centrifuged for 10 min at 12,000 rpm.

The supernatant was removed. For washing, an amount of 500 µl 70% ethyl alcohol was added and centrifuged for 8 min at 12,000 rpm. The supernatant was decanted or tipped and the tubes blotted on Whatman paper or clean tissue for 15 min. For apoptosis, once the tube was seen to be dry, the pellet was resuspended in 50 µl or appropriate volume of TE buffer (10 mM tris, 1mM EDTA, pH8), and supplemented with 5% glycerol for 30 min. The resuspended DNA with 6 x loading buffer was loaded directly on gel. So, the fine apoptotic bands could be detected even in the control sample.

#### **Apoptosis analysis:**

For apoptosis, the extracted DNA was gently resuspended with TE buffer supplemented with 5% glycerol, gently pipetted, and then mixed with 6 x loading buffer, and loaded directly on the gel.<sup>21</sup> The remaining DNA was kept at -20°C for another loading. Apoptotic bands appeared and located at 180 bp.

#### **Biochemical analysis:**

Serum total protein, cholesterol, triglycerides, glucose, ALT, AST, GGT, ALP and LDH levels were determined automatically using Integra 800 auto-analyzer, National Liver Institute, Menoufiya University. Lead was determined in serum samples by atomic absorption spectrophotometry using G.B.C. 902, double beam atomic absorption spectrophotometer (A.D.S. instrument), Faculty of Agriculture, Menoufiya University.

#### **Statistical analysis:**

Data were represented in tabular form as mean ± standard deviation. Students t-test was used for evaluating statistical significance for results according to **Hine and Wetherill**.<sup>22</sup>

## **Results**

#### **Body weight:**

Decreased significantly by 33.83% after heat stress for 25 days and by 41.61% after lead treatment under hyperthermia. Vitamin supplementation reduced the effect of heat and lead on body weight.

#### **Hepatosomatic index:**

Heat stress induced elevation in hepatosomatic index (liver (g)/body weight(g) ratio) by 15.68%; however lead under hyperthermia aggravated this effect to 20.41%. On the other hand treatment with antioxidants with rehydration ameliorated these effects.

#### **Biochemistry:**

Heat stress resulted in significant increase in serum glucose, cholesterol, triglycerides, ALT and LDH levels. On the other hand, serum ALP activity decreased significantly. However serum total protein, lead, AST and GGT levels were not affected.

Lead induced significant increase in serum glucose, total protein, lead, ALT, AST, GGT, ALP and LDH levels under heat stress conditions. However serum cholesterol decreased significantly and triglycerides were not significantly affected.

Treatment of lead toxicity under heat stress conditions with rehydran-n solution and vitamins (C & E), silymarin, and selenium supplements improved all biochemical parameters under investigation, especially total protein, AST, GGT and ALP levels.

#### **Molecular findings:**

Heat stress induced fine apoptotic bands indicating high degree of liver cell destruction. Lead induced aggravation of apoptosis in liver cells under heat stress conditions. Treatment with vitamins (C & E), silymarin, and rehydration solution ameliorated these effects.

#### **Histopathology:**

Under heat stress conditions, randomly scattered hepatocytes showed acidophilic and apoptotic changes. Rats exposed to lead and heat stress showed marked acidophilic and apoptotic changes in zone 2 (mid acinar zone) with microvesicular steatosis in zone 1 (periportal areas).

Vitamins C and E and other supplements resulted in significant reduction of both lead, and to some extent heat stress manifestations.

### Discussion

Heat is a stressor that evokes several physiological reactions in humans and animals.<sup>23</sup> Many biochemical and physiological systems of the body are affected by exposure to heat, such as enzymatic, metabolic, cardiovascular, respiratory, haematopoietic, endocrinal and immunological systems, as well as blood and body fluid composition.<sup>24-33</sup> Lead intake causes toxic effects on different body systems.<sup>34</sup> Hyperthermia exaggerates the toxic effects of lead. There is evidence that combined supplementation with vitamin C and silymarin ameliorates the toxic effects of lead under mild conditions.

The present study showed that heat stress resulted in significant elevations in blood glucose, cholesterol, triglycerides, ALT and LDH levels.

**Mertsching** showed that the decrease of insulin release in heat stressed animals decreases the rate of glucose utilization in tissues which lead to increased blood glucose level.<sup>35</sup> Heat stress causes disturbances in liver function tests in hamsters,<sup>36</sup> rats,<sup>37</sup> and humans.<sup>38</sup> In humans and rats exposed to hyperthermia, autophagic vacuolation, and dilatation of both Golgi apparatus and endoplasmic reticulum, together with elevation of serum GOT, GPT and LDH activities, have been reported.

Our results indicated that lead induced significant elevations of blood glucose, total protein, lead, ALT, AST, GGT, ALP and LDH levels under heat stress conditions. These results are in agreement with those reported by earlier investigators (*vide supra*).

Hyperthermia was reported to induce RBCs hemolysis,<sup>39</sup> which may be responsible for increased serum total protein. **Latner** showed that the increase in serum GGT is an indicator of liver fibrosis and correlates with the development of hepatobiliary diseases indicating toxic liver damage.<sup>40</sup>

**Eissa et al** reported increased serum alkaline phosphatase activity in workers subjected to lead.<sup>41</sup> This was interpreted to changes in membrane permeability<sup>42</sup> that lead to hypoxia of hepatocytes<sup>43</sup> and increased peroxidation.<sup>44</sup> The significant increase in serum ALT

and AST activities in lead-intoxicated rats under heat stress conditions reflects the damage of cells and alterations in cell permeability.<sup>45</sup> In accordance with our findings, **Skoczynska et al** showed a decrease in plasma cholesterol in rats treated with small doses of lead.<sup>46</sup>

Recent studies have shown that lead induces hepatic DNA damage.<sup>47-49</sup> In the present study lead aggravated apoptotic DNA fragmentation induced by heat stress. **Pagliara et al** showed that lead induced hepatic hyperplasia followed by apoptosis mediated by oxidative stress in kupffer cells.<sup>50</sup>

The results of this study showed decreased body weight in both heat stressed groups. Decrease in body weight due to hyperthermia may be attributed to alterations of the availability of enzyme substrates and hormones, and to the decrease of food intake and excessive water loss<sup>51-54</sup> It may be related to increased catabolism and tissue destruction under heat stress conditions.

The loss of body weight is generally considered to be a physiological index of lead intoxication. **Gerber et al**<sup>55</sup> and **Abou-El Maged et al**<sup>56</sup> attributed the decrease of body weight to strong affinity of lead for -SH, amine, carboxylic and phosphate ligand groups of biological membranes and protein. This leads to enzyme inhibition and disruption of numerous metabolic pathways including those of oxidative phosphorylation<sup>57</sup> causing general inhibition of metabolism, besides other factors such as malabsorption of food materials, depression of appetite, and reduction of food consumption.<sup>58</sup> Further, some authors attribute the growth retardation to hormonal imbalance due to lead toxicity.<sup>59-61</sup>

Our results show that heat aggravates hepatomegaly, and lead pronounces this effect. This may be due to the diffuse wide spread affection of zone 1 (periportal areas) by microvesicular steatosis, and the mild mononuclear inflammatory cell infiltration.

Zone 1 is upstream in the blood flowing to the liver acinus, and zone 3 is downstream. Therefore zone 1 hepatocytes are exposed to the highest concentration of lead compared to zone 3 hepatocytes. The periportal distribution of lead-induced microvesicular steatosis indicates a direct hepatotoxic effect of lead, and not through metabolic byproducts of lead.

It has been shown that lead nitrate induces a synchronized wave of hepatocyte proliferation in rat liver.<sup>62</sup> **Pagliara et al** showed that lead induces liver hyperplasia followed by apoptosis mediated by oxidative stress in Kupffer cells.<sup>63</sup>

Vitamin E is a known antioxidant that protect against toxicity through its free radical scavenging capacity; therefore, it can prevent lipid peroxidation and fix the biomembranes.<sup>64</sup> It can also prevent membrane phospholipid degradation through phospholipidases linkage.<sup>65</sup> Vitamin C has been reported to reduce the possibility of lead interacting with critical biomolecules and thus preventing its toxicity. It can be used as a chelator for lead to decrease the risk of its toxic effects.<sup>66</sup> The preventive activity of vitamin C may be related to its antioxidant efficacy that inhibits lipid peroxidation enhanced by lead. **Blankenship et al** showed that vitamin C protected cells from undergoing apoptosis.<sup>67</sup>

Silymarin has anti-inflammatory activities mediated by alteration of Kupffer cell function.<sup>68</sup> It has been reported that silymarin improved liver function tests related to hepatocellular necrosis and/or increased membrane permeability.<sup>69</sup> Its protective effect is attributed to its antioxidant and free radicals scavenging properties. Silymarin has been shown to reverse histopathological changes of CCl<sub>4</sub>, such as necrosis, fatty change, ballooning degeneration, and inflammatory infiltration of lymphocytes around the central vein. In agreement of our results silymarin was found to reduce hepatic collagen accumulation by 35% in rats with secondary biliary cirrhosis.<sup>70</sup> **Saravanan et al** showed that vitamin C or/and silymarin were hepatoprotective and have antioxidant effect against ethanol intoxication.<sup>71</sup> The hepatoprotective effect of silymarin may be attributed to its ability to scavenge oxygen free radicals, and inhibition of liver mitochondria lipid peroxidation.

The increase of water intake is an important means in amelioration of heat stress effects. Rehydran-n solution has high water content in addition to sodium, potassium, and glucose that improve the osmotic fragility of cellular membranes and hence may ameliorate heat stress.

In conclusion, combined supplementation with vitamins C, E, selenium, silymarin, and rehydran-n solution markedly reduce the hepatotoxic effects of lead under heat stress conditions, and can be used as a protective measure in such circumstances.

## REFERENCES

1. Chia KS, Jeyaratanam J, Tan C, Ong HY, Ong CN, Lee E. Glomerular function of lead-exposed workers. *Toxicol lett.* 1995; 77: 319-328.
2. Sandhir R, Gill KD. Effect of lead on lipid peroxidation in liver of rats. *Biol Trace Elem Res.* 1995; 48: 91-97.
3. Upasani CD, Khera A, Balaraman R. Effect of lead with vitamins E, C, or spirulina on malondialdehyde: Conjugated dienes and hydroperoxides in rats. *Indian J Exp Biol.* 2001; 39: 70-74.
4. Fracasso ME, Perbellini L, Solda S, Talamini G, Franceschetti P. Lead induced DNA strand breaks in lymphocytes of exposed workers: role of reactive oxygen species and protein kinase C. *Mutat Res.* 2002; 515: 159-169.
5. Valverde M, Fortoul TE, Diaz-Barriga F, Mejia J, del Castillo ER. Genotoxicity induced in CD-1 mice by inhaled lead in differential organ response. *Mutagenesis.* 2002; 17: 55-61.
6. Shalan MG, Mostafa MS, Hassouna MM, Hassab el-Nabi SE, El-Refaie A. Amelioration of lead toxicity on rat liver with vitamin C and silymarin supplements. *Toxicology.* 2005; 206: 1-15.
7. Honchel R, Marsanco L, Cohen D, Shedlofsky S, McClain CJ. Lead enhances lipopolysaccharide and tumor necrosis factor liver injury. *J Lab Clin Med.* 1991; 117: 202-208.
8. Pagliara P, Carla EC, Caforio S, Chionna A, Massa S, Abbro L, Dini L. Kupffer cells promote lead nitrate-induced hepatocyte apoptosis via oxidative stress. *Comp Hepatol.* 2003 b; 2: 8.
9. Kamal TH, Bahgat MM, Abo-Grween LGA. Effect of heat on blood components, prothrombin time and growth hormone in immature male rats. *J Biomed Sci Ther.* 1992b; 8: 50-58.
10. Weniger JH, Methias J, Forster E. Influence of temperature and humidity on growth. *J Anim Breed Genet.* 1991; 108: 379-388.
11. Tharwat EE, Amin SO, Khadr AF, Kotby EA. The physiological response of New Zealand white rabbit to heat stress. *Cah Options Mediterr.* 1994; 8: 634-637.
12. Groenink K, Van-Der-Gugten J, Zethof-T, Van-Der-Heyden J, Olivier B. Stress-induced hyperthermia in mice: Hormonal correlates. *Physiol Behav.* 1994; 56: 747-749.
13. Michael MI. Effect of lead on some physiological parameters in heat stressed rats. 1997; MSc. Thesis, Institute of Studies and Environmental research, Ain Shams University, Egypt.



14. Llobet JM, Dominco JL, Paternain JL, Crobell J. Treatment of acute intoxication. A quantitative comparison of a number of chelating agents. *Arch Environ Contam Toxicol.* 1990; 19: 185-189.
15. Hsu PC, Guo YL. Antioxidant nutrients and lead toxicity. *Toxicology.* 2002; 180: 33-44.
16. Shalan MG. Effect of radiation on some biochemical parameters and the prophylactic action of some chemical treatment. 2000; PhD Thesis, Faculty of Science, Menoufiya University, Egypt.
17. Soto CP, Perez BL, Favari LP, Reyes JL. Prevention of alloxan-induced diabetes mellitus in the rat by silymarin. *Comp Biochem Physiol Pharmacol Toxicol Endocrinol.* 1998; 119: 125-129.
18. Hagymasi K, Kocsis I, Lugasi A, Fesher J, Blazovics A. Extrahepatic biliary obstruction: Can silymarin protect liver function? *Phytother Res.* 2002; 16 (suppl. 1): s78-s80.
19. Lucena MI, Andrade RJ, de la Cruz JP, Rodriguez-Mendizabal M, Blanco R, de la Cuesta S. Effects of silymarin MZ-80 on oxidative stress in patients with alcoholic cirrhosis. Results of randomized, double-blind, placebo-controlled clinical study. *Int J Clin Pharmacol Ther.* 2002; 401: 2-8.
20. Aljanabi SM, Martinez I. Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. *Nucl Acids Res.* 1997; 25: 4692-4693.
21. Hassab El-Nabi SE. Molecular and cytogenetic studies on the antimutagenic potential of eugenol in human lymphocytes culture treated with depakine and apetryl drugs. *J Egypt Ger Soc Zool.* 2004 ; 43: 171-196.
22. Hine J, Wetherill GB. The t-test X2 Goodness of fit. In: *A Programmed Test in Statistics. Book 3.* 1975. Chapman and Hill, London.
23. Amer MM. Lead and lead chelation in relation to physiological and histological aspects of heat stressed rats. 1997. PhD Thesis, Faculty of Science, Ain Shams University, Egypt.
24. Adolph EF. *Physiology of Man in the Desert.* 1947. Intersci Pub Inc. New York.
25. Bass DE, Kleenan CR, Quinn M, Henshel A, Hegnavar AH. Mechanism of acclimatization to heat in men. *Med.* 1955; 34: 323-325.
26. UNESCO. Environmental physiology and psychology in arid conditions. *Proc Lucknow Symp.* 1964. Arid Zone Research. UNESCO, place de Fontenoy, Paris 7e XXIT: 1-400.
27. Kamal TH. Physiological reactions of cows to hot environmental conditions. In: *Radioisotopes in Animal Nutrition and Physiology (Proc. Symp. Prague, Czechoslovakia, Nov. 23-27, 1964).* Proc. Series IAEA, Vienna: 767-792.
28. Francesconi RD, Hubbard RW, Mager M. Chronic low-sodium diet in rats: hormonal and physiological effects during exercise in the heat. *J Appl Physiol Res Environ Exercise Physiol.* 1983a; 55: 870-874.
29. Francesconi RD, Sauvka MN, Pandoll RB. Hypohydration and heat acclimation: plasma renin and aldosterone during exercise. *J Appl Physiol.* 1983 b; 55: 1790-1794.
30. Kamal TH, Bahgat MM. Effect of heat on the pattern of distribution of immunoglobulins. *Egypt Med Assoc.* 1987; 70: 275-281.
31. Kamal TH, Bahgat MM, Abo-Grween LGA. Heat as immuno-suppressive agent on the in vivo production of antibodies in immature male rats. *J Egypt Med Assoc.* 1992 a; 75: 99-104.
32. Ibrahim MS, Kamal TH, Mostafa SI, Mostafa AI. Individual variation among Egyptian men in heat tolerance. *J Egypt Med Assoc.* 1992; 72: 91-98.
33. Kamal TH. Indices of heat tolerance and amelioration of heat stress. In: *Prospects of buffalo production in the Mediterranean and Middle East. Proc Jt ESAP, IAAP, FAO, ICAMAS & OIE Symp, Cairo, 9-12 Nov, 1992.* Pudoc Sci Pubs. Wageningen, PP. 198-200.
34. Mahoffey KR, Michaelson IA. The interaction between lead and nutrition. In: Needleman HL (ed). *Low Level Lead Exposure: The Clinical Implication of Current Research.* 1980. New York: Raven. 159-200.
35. Mertsching HJ. The effects of high environmental temperature on prolactin and glucose in fed and fasted monolactating dairy cattle. MSc Thesis, 1981. University of Missouri, Columbia.
36. Ilana I, Reuben C, Yaircassuto. Energy metabolism in kidney of heat acclimated hamsters. *Am J Physiol.* 1975; 229: 1234-1236.
37. Hassanin SH, Khalil EA, Abd-El Aziz AMS, El-Sobhy H.E. Changes in some physiological parameters of albino rats at different ambient temperatures. *Asian-Austral J Anim Sci.* 1994; 7: 471-474.
38. Benjamin SA, Nikulo KJ, Powers BE, Haden FF. Radiation and heat. In: Hasckek WM, Rousseaux CG (eds). *Handbook of Toxicologic Pathology.* 1991. Academic Press Inc, USA. 1042-1043.

39. Kimber RJ, Lander H. The effect of heat on human red cell morphology, fragility and subsequent survival in vivo. *J Lab Clin Med.* 1964; 64: 922.
40. Latner AL. *Clinical Biochemistry.* 7<sup>th</sup> edn, 1975. WB Saunders Company, Philadelphia, London, Toronto.
41. Eissa SA, El-Sayed EA, Mohamed HA, Marrie M. Liver function among workers exposed to lead in battery industry. *J Egypt Ger Soc Zool.* 1993; 10: 373-383.
42. Steffensen IL, Mesna OJ, Andruchow E, Namork E, Hyltand K, Andersen RA. Cytotoxicity and accumulation of Hg, Ag, Cd, Cu, Pb and Zn in human peripheral T and B lymphocytes and monocytes in vitro. *Gen Pharmacol.* 1994; 25: 1621-1633.
43. Zakaria YA, El-Batanouni MM, Amer FB, Dessoaky SA. Hepatic dysfunction among workers exposed to arsenic and lead glass industry. *Med J Cairo Univ.* 1986; 53: 1.
44. Skozynska A, Smolik A. The effect of combined exposure to lead and cadmium on serum lipids and lipid peroxides level in rats. *Int J Occup Med Environ Health.* 1994; 7: 263-271.
45. Harvey AM, Jons RJ, McKsick VA, Owens AH, Ross RS. Normal and abnormal hepatic physiology. In: *Principles and Practice of Medicine.* 20<sup>th</sup> edn, 1980; Appleton Century Crofts, Prentice-Hall Inc. 504-702.
46. Skoczynska A, Smolik R, Jelen M. Lipid abnormalities in rats given small doses of lead. *Arch Toxicol.* 1993; 67: 200-204.
47. Danadevi K, Rozati R, Sleha-Banu B, Hanumanth Rao P, Grover P. DNA damage in workers exposed to lead using Comet assay. *Toxicology.* 2003; 187: 183-193.
48. Hengstler JG, Bolm-androff U, Faldum A, Janssen K, Reifen-rath M, Gotte W, et al. Occupational exposure to heavy metal: DNA damage induction and DNA repair inhibition prove co-exposures to cadmium, cobalt and lead as more dangerous than hitherto expected. *Carcinogenesis.* 2003; 24: 63-73.
49. Xu DX, Shen HM, Ahu QX, Chua CN. The association among semen quality, oxidative DNA damage in human spermatozoa and concentrations of cadmium, lead and selenium in seminal plasma. *Mutat Res.* 2003; 534: 155-163.
50. Pagliara P, Chionna A, Carla EC, Caforio S, Dini L. Lead nitrate and gadolinium chloride administration modify hepatocyte cell surfaces. *Cell Tiss Res.* 2003 a; 312: 41-48.
51. Kamal TH. Thermal effects of various temperature humidity combinations on Holstein cattle as measured by eight physiological responses. *Missouri Agr Res Bul.* 1964; 863.
52. Hafez ESE. Environmental effects on animal productivity. In: *Adaptation of Domestic Animals.* 1968. Lea & Febiger, Philadelphia. 74-96
53. Lofgreen GP, Givens RC, Marrison SR. Effect of drinking water temperature on beef cattle performance. *J Anim Sci.* 1975; 40: 322.
54. Collier RJ, Beed DK, Thacher WW, Israel LA, Wilcox CJ. Influence of environment and its modification on dairy animals health and production. *J Dairy Sci.* 1982; 56: 2213-2227.
55. Gerber GB, Leonard A, Jacquet P. Toxicity, mutagenicity and tetratogenicity of lead. *Mut Res.* 1980; 76: 145.
56. Abo-El Maged A, Shore AH, Abd El-Ghany SM, El-Deeb TS. Some biochemical indices of lead and cadmium exposure in healthy and protein energy malnourished children in Assiut locality. *J Union Arab Biol.* 1994; 1: 437-452.
57. Bryce-Smith D, Stephens R. Sources and effects of environmental lead. In: Ros J (ed). *Trace Elements in Health.* 1983. Butterworth Mid-Country Press, London. 83.
58. Hammond PB, Minnema DJ, Shuulka R. Lead exposure lowers the set point for food consumption and growth in weaning rats. *Toxicol Appl Pharmacol.* 1990; 106: 80-87.
59. Huseman CA, Moriarty CM, Angle CR. Childhood lead toxicity and impaired release of thyrotropin-stimulating hormone. *Environ Res.* 1987; 42: 524-532.
60. Angle CR, Kuntzelman DR. Increased erythrocyte protoporphyrins and blood lead. A pilot study of childhood growth patterns. *J Toxicol Environ Health.* 1989; 26: 149-156.
61. Camoratto AM, White LM, Lau YS, Ware GO, Berry WD, Moriarty CM. Effect of exposure to low level lead on growth and growth hormone release in rats. *Toxicol.* 1993; 83: 101-114.
62. Kubo Y, Younaga M, Masuttara M, Terai S, Nakamura T, Okita K. Hepatocyte proliferation induced in rats by lead nitrate is suppressed by several tumor necrosis factor inhibitors. *Hepatology.* 1996; 23: 104-114.
63. Pagliara P, Carla EC, Caforio S, Chionna A, Massa S, Abbro L, Dini L. Kupffer cells promote lead nitrate-induced hepatocyte apoptosis via oxidative stress. *Comp Hepatol.* 2003 b; 2: 8.

64. Upasani CD, Khera A, Balaraman R. Effect of lead with vitamins E, C, or spirulina on malondialdehyde: Conjugated dienes and hydroperoxides in rats. *Indian J Exp Biol.* 2001; 39: 70-74.
65. Konings AWT, Drijver EB. Radiation effects in membranes. I. Vitamin E deficiency and lipid peroxidation. *Rad Res.* 1979; 80: 494-501.
66. Simon JA, Hudes ES. Relationship of ascorbic acid to blood lead levels. *JAMA.* 1999; 281: 2289-2293.
67. Blankenship LJ, Claiste DL, Wise JP, Orenstein JM, Dye LE, Patierno SR. Induction of apoptotic cell death by particulate lead chromate: differential effects of vitamin C and E on genotoxicity and survival. *Toxicol Appl Pharmacol.* 1997; 146: 270-280.
68. Dehmlo WC, Erhard J, DeGroot II. Inhibition of kupffer-cell function as an explanation for the hepatoprotective properties of silibinin. *Hepatology.* 1996; 23: 749-754.
69. Buzzelli G, Mosarella S, Giusti A, Duchini AP, Morena C, Lampertio M. A pilot study on the liver. Protective effect of silybin-phosphatidyl choline complex (IDB 1016) in chronic active hepatitis. *Int J Clin Pharmacol Ther Toxicol.* 1993; 31: 456-460.
70. Jia JD, Bauer M, Cho JJ, Ruchl M, Milani S, Boigk G, et al. Antifibrotic effect of silymarin secondary biliary fibrosis is mediated by down regulation of procollagen alpha 1 (I) and TIMP-1. *J Hepatol.* 2001; 35: 392-398.
71. Saravanan R, Prakasam A, Ramesh B, Pugalendi KV. Influence of piper beetle on hepatic marker enzymes and tissue antioxidant status in ethanol-treated Wistar rats. *J Med Food.* 2002; 5: 197-204.